

UT-945

June, 2001

## Warped Compactification with a Four-Brane

S. Hayakawa and Izawa K.-I.

*Department of Physics, University of Tokyo,  
Tokyo 113-0033, Japan*

### Abstract

Warped compactification of the six-dimensional bulk with a negative cosmological constant is realized with a 4-brane along with an abelian gauge theory. No fine tuning of couplings are needed to obtain the vanishing cosmological constant in four dimensions, as is the case for the bulk with a positive cosmological constant.

# 1 Introduction

Higher-dimensional approach to the cosmological constant problem considered by Rubakov and Shaposhnikov [1] is based on the idea that the vanishing cosmological constant in four dimensions might be achieved through warped compactification even in the presence of a nonvanishing higher-dimensional cosmological constant. This might be possible since effective four-dimensional flatness is not necessarily in contradiction with highly curved spacetime as a higher-dimensional space.

In a previous paper [2], we investigated the case of the six-dimensional bulk with a positive cosmological constant and achieved the vanishing four-dimensional cosmological constant without metric singularity by introducing an abelian gauge field [3]. In this paper, we proceed to the case of a negative bulk cosmological constant to get regular and compact extra dimensions. We are led to introduce a 4-brane in the bulk [4] along with an abelian gauge field, which is common to the case for a positive cosmological constant.

## 2 The Model

Let us assume an  $SO(2)$ -symmetric warped metric as a six-dimensional background [1]-[4]:

$$ds^2 = g_{MN}dx^M dx^N = \sigma(r)\bar{g}_{\mu\nu}dx^\mu dx^\nu - dr^2 - \rho(r)d\theta^2, \quad (1)$$

where  $\bar{g}_{\mu\nu}$  denote the four-dimensional metric independent of  $(r, \theta)$  with  $0 \leq \theta < 2\pi$ . The action of the metric and an abelian gauge field with a 4-brane at  $r = r_1$  is given by

$$S = \int d^6x \sqrt{-g} \left( \frac{1}{2}R - \frac{1}{4}F_{MN}F^{MN} - \Lambda \right) - \int d^6x \sqrt{-g_5} \lambda \delta(r - r_1), \quad (2)$$

where  $\Lambda < 0$  and  $g_5$  denotes the determinant of the induced metric on the 4-brane.

The gauge field equations of motion are obtained as

$$\partial_M(\sqrt{-g}F^{MN}) = 0. \quad (3)$$

The background configuration with the  $SO(2)$  symmetry is given by

$$A_\mu = A_r = 0, \quad A_\theta = a(r), \quad (4)$$

which yield the following field strength:

$$F^{r\theta} = \frac{B}{\sigma^2 \sqrt{\rho}}, \quad (5)$$

where  $B$  is an integration constant. With this gauge field configuration, the Einstein equations lead to

$$\frac{3}{2} \frac{\sigma''}{\sigma} + \frac{3}{4} \frac{\sigma' \rho'}{\sigma \rho} - \frac{1}{4} \frac{\rho'^2}{\rho^2} + \frac{1}{2} \frac{\rho''}{\rho} - \frac{\Lambda_4}{\sigma} = -\frac{B^2}{2\sigma^4} - \Lambda - \lambda\delta(r - r_1), \quad (6)$$

$$\frac{3}{2} \frac{\sigma'^2}{\sigma^2} + \frac{\sigma' \rho'}{\sigma \rho} - \frac{2\Lambda_4}{\sigma} = \frac{B^2}{2\sigma^4} - \Lambda, \quad (7)$$

$$2 \frac{\sigma''}{\sigma} + \frac{1}{2} \frac{\sigma'^2}{\sigma^2} - \frac{2\Lambda_4}{\sigma} = \frac{B^2}{2\sigma^4} - \Lambda - \lambda\delta(r - r_1), \quad (8)$$

where the prime denotes differentiation with respect to  $r$  and we have used four-dimensional Einstein equations for the metric  $\bar{g}_{\mu\nu}$  with the cosmological constant  $\Lambda_4$ , which comes out as an integration constant.

We get the junction conditions across the 4-brane from the equations of motion:

$$\sigma'(r_1 + 0) - \sigma'(r_1 - 0) = -\frac{1}{2}\lambda\sigma(r_1), \quad (9)$$

$$\rho'(r_1 + 0) - \rho'(r_1 - 0) = -\frac{1}{2}\lambda\rho(r_1). \quad (10)$$

In the regions  $r < r_1$  and  $r > r_1$ , the bulk Einstein equations are reduced to an equation of motion

$$z'' = -\frac{\partial V(z)}{\partial z}; \quad V(z) = \frac{25}{96}B^2z^{-6/5} + \frac{5}{16}\Lambda z^2 - \frac{25}{24}\Lambda_4z^{6/5}, \quad (11)$$

$$\sigma = z^{4/5}, \quad \rho = C_{\mp}^{-2}z'^2z^{-6/5}, \quad (12)$$

where  $C_{\mp}$  denote an integration constant for each region. Note that this equation describes the motion of a particle with the position  $z$  at the time  $r$  in a potential  $V(z)$ .

### 3 The Solutions

Now we seek regular metric solutions in the desirable case of  $\Lambda_4 = 0$ . The junction conditions Eqs.(9) and (10) can be expressed in terms of  $z$  under  $\lambda \neq 0$ :

$$z'_+ - z'_- = -\frac{5}{8}\lambda z_1, \quad (13)$$

$$z'_+z'_- = -\frac{5}{16}B^2z_1^{-6/5} + \frac{5}{8}\Lambda z_1^2, \quad (14)$$

where we have denoted  $z'_{\pm} = z'(r_1 \pm 0)$  and  $z_1 = z(r_1)$  (which may be normalized to one without loss of generality). Eqs.(13) and (14) can be solved for  $z'_{\pm}$ , provided  $5\lambda^2 + 32\Lambda \geq 0$ , with  $z'_- > 0$  and  $z'_+ < 0$ . This change of the velocity  $z'$  from positive to negative is crucial for compactification of the extra dimensions, which is achieved by means of the 4-brane with  $\lambda > 0$ .

The equation of motion Eq.(11) implies  $z' = 0$  at  $r = r_0, \bar{r}$  with  $z' \neq 0$  for  $r_0 < r < r_1$  and  $r_1 < r < \bar{r}$ . The ‘energy’ conservations as the motion of the position  $z$  in the ‘time’ regions  $r_0 \leq r < r_1$  and  $r_1 < r \leq \bar{r}$  yield

$$\frac{25}{96}B^2z_0^{-6/5} + \frac{5}{16}\Lambda z_0^2 = \frac{1}{2}z_-'^2 + \frac{25}{96}B^2z_1^{-6/5} + \frac{5}{16}\Lambda z_1^2, \quad (15)$$

$$\frac{25}{96}B^2\bar{z}^{-6/5} + \frac{5}{16}\Lambda \bar{z}^2 = \frac{1}{2}z_+'^2 + \frac{25}{96}B^2z_1^{-6/5} + \frac{5}{16}\Lambda z_1^2, \quad (16)$$

respectively. Here we have denoted  $z_0 = z(r_0)$ ,  $\bar{z} = z(\bar{r})$ , whose values are determined by these equations.

Continuity of  $\rho$  across the 4-brane imposes a condition

$$C_-^{-2} z_-'^2 = C_+^{-2} z_+'^2 \quad (17)$$

from Eq.(12). For regularity of the higher-dimensional metric, we have conditions

$$(\sqrt{\rho})'(r_0) = -(\sqrt{\rho})'(\bar{r}) = 1, \quad (18)$$

that is,

$$C_- z_0^{3/5} = \frac{5}{16} B^2 z_0^{-11/5} - \frac{5}{8} \Lambda z_0, \quad (19)$$

$$C_+ \bar{z}^{3/5} = \frac{5}{16} B^2 \bar{z}^{-11/5} - \frac{5}{8} \Lambda \bar{z}. \quad (20)$$

The above three conditions can be solved for three integration constants  $B$  and  $C_{\mp}$ , which yield a reflection symmetric metric with respect to the 4-brane.

Namely, the effective four-dimensional cosmological constant  $\Lambda_4$  can vanish for generic values of  $\Lambda < 0$  and  $\lambda > 0$  with the extra dimensions compactified.

We note that, in addition to the 4-brane at  $r = r_1$ , 3-branes may be introduced at  $r = r_0$  and/or  $r = \bar{r}$  straightforwardly. For example, if we put a 3-brane with the tension  $\lambda_0$  at  $r = r_0$ , the regularity condition Eq.(19) is replaced by [2]

$$C_- (1 - \frac{\lambda_0}{2\pi}) z_0^{3/5} = \frac{5}{16} B^2 z_0^{-11/5} - \frac{5}{8} \Lambda z_0, \quad (21)$$

which also allows the full equations of motion to be satisfied without singularity.

## 4 Conclusion

We have obtained the regular metric with compact extra dimensions in the case of the negative bulk cosmological constant in six dimensions. The four-dimensional cosmological constant  $\Lambda_4$  can vanish without fine tuning of Lagrangian parameters  $\Lambda < 0, \lambda > 0$  (and the tensions of 3-branes, if present).

We note that regular and compact metrics are also present under  $\Lambda_4 \neq 0$ , as is the case for a positive bulk cosmological constant [2]. Namely,  $\Lambda_4$  is none other than an integration constant to be determined by boundary conditions, which might be supplied by some other sector to be added in the model [5].

## References

- [1] V.A. Rubakov and M.E. Shaposhnikov, Phys. Lett. **B125** (1983) 139.
- [2] S. Hayakawa and Izawa K.-I., Phys. Lett. **B493** (2000) 380 [hep-th/0008111].
- [3] C. Wetterich, Nucl. Phys. **B255** (1985) 480;  
G.W. Gibbons and D.L. Wiltshire, Nucl. Phys. **B287** (1987) 717.
- [4] Z. Chacko and A.E. Nelson, Phys. Rev. **D62** (2000) 085006 [hep-th/9912186].
- [5] Izawa K.-I., Prog. Theor. Phys. **105** (2001) 185 [hep-ph/0007079].